

KASHAEV'S CONJECTURE AND THE CHERN–SIMONS INVARIANTS OF KNOTS AND LINKS

HITOSHI MURAKAMI, JUN MURAKAMI, MIYUKI OKAMOTO, TOSHIE TAKATA,
AND YOSHIYUKI YOKOTA

ABSTRACT. R.M. Kashaev conjectured that the asymptotic behavior of his link invariant, which equals the colored Jones polynomial evaluated at a root of unity, determines the hyperbolic volume of any hyperbolic link complement. We observe numerically that for knots 6_3 , 8_9 and 8_{20} and for the Whitehead link, the colored Jones polynomials are related to the hyperbolic volumes and the Chern–Simons invariants and propose a complexification of Kashaev's conjecture.

1. INTRODUCTION

In [5], R.M. Kashaev defined a link invariant associated with the quantum dilogarithm, depending on a positive integer N , which is denoted by $\langle L \rangle_N$ for a link L . Moreover, in [6], he conjectured that for any hyperbolic link L , the asymptotics at $N \rightarrow \infty$ of $|\langle L \rangle_N|$ gives its volume, that is

$$\text{vol}(L) = 2\pi \lim_{N \rightarrow \infty} \frac{\log |\langle L \rangle_N|}{N}$$

with $\text{vol}(L)$ the hyperbolic volume of the complement of L . He showed that this conjecture is true for three doubled knots 4_1 , 5_2 , and 6_1 . Unfortunately his proof is not mathematically rigorous.

Afterwards, in [9], the first two authors proved that for any link L , Kashaev's invariant $\langle L \rangle_N$ is equal to the colored Jones polynomial evaluated at $\exp(2\pi\sqrt{-1}/N)$, which is written by $J_N(L)$, and extended Kashaev's conjecture as follows.

Conjecture 1.1 (Volume Conjecture).

$$\|L\| = \frac{2\pi}{v_3} \lim_{N \rightarrow \infty} \frac{\log |J_N(L)|}{N},$$

where $\|L\|$ is the simplicial volume of the complement of L and v_3 is the volume of the ideal regular tetrahedron.

Note that the hyperbolic volume $\text{vol}(L)$ of a hyperbolic link L is equal to $\|L\|$ multiplied by v_3 . This conjecture is not true for links in general, as $J_N(L)$ vanishes for a split link L . Note also that it is shown by Kashaev and O. Tirkkonen in [4] that the volume conjecture holds for torus knots. See [11] and [14, 13] for discussions about Kashaev's conjecture for hyperbolic knots from the viewpoint of tetrahedron decomposition.

In this paper, following Kashaev's way to analyze the asymptotic behavior of the invariant, we observe numerically, by using MAPLE V (a product of Waterloo Maple Inc.) and SnapPea [12], that for the hyperbolic knots 6_3 , 8_9 , 8_{20} , and for the Whitehead link, the colored Jones

This research is partially supported by Grand-in-Aid for Scientific Research, The Ministry of Education, Science, Sports and Culture.

polynomials are related to the hyperbolic volumes and the Chern–Simons invariants. Note that the knots 6_3 and 8_9 are not doubles of the unknot.

We also discuss a relation between the asymptotic behavior of $J_N(L)$ and the Chern–Simons invariant of the complement of the above-mentioned links L , and propose the following conjecture.

Conjecture 1.2 (Complexification of Kashaev’s conjecture). *Let L be a hyperbolic link. Then the following formula holds.*

$$J_N(L) \sim \exp \frac{N}{2\pi} (\text{vol}(L) + \sqrt{-1} \text{CS}(L)) \quad (N \rightarrow \infty)$$

where $\text{CS}(L)$ is the Chern–Simons invariant of L [1, 8]. Note that the complement of L is a hyperbolic manifold with cusps.

The statement of this conjecture will be given more properly in the last section.

Acknowledgments. We thank the participants in the meeting “Volume conjecture”, October 1999 and those in the workshop “Recent Progress toward the Volume Conjecture”, March 2000, both of which were held at the International Institute for Advanced Study. The latter was financially supported by the Research Institute for Mathematical Sciences, Kyoto University. We are grateful to both of the institutes.

H.M., J.M. and M.O. express their gratitude to Graduate School of Mathematics, Kyushu University, where the essential part of this work was carried out in December 1999.

Thanks are also due to S. Kojima for his suggestion of the Chern–Simons invariant, to K. Mimachi for valuable discussions, and to K. Hikami for informing Pari-Gp [2] and fitting.

2. PRELIMINARIES

First we will briefly review the colored Jones polynomials of links following [7]. It is obtained from the quantum group $U_q(\mathfrak{sl}(2, \mathbb{C}))$ and its N -dimensional irreducible representation.

Let L be an oriented link. We consider a $(1, 1)$ -tangle presentation of L , obtained by cutting a component of the link. We assume that all crossing and local extreme points are as in Figure 1. We can calculate the N -colored Jones polynomial $J_L(N)$ evaluated at the N -th root of unity for L in the following way. We start with a labeling of the edges of the $(1, 1)$ -tangle presentation with labels $\{0, 1, \dots, N-1\}$. Here we label the two edges containing the end points of the tangle by 0. Following the labeling, we associate a positive (respectively negative) crossing with the element R_{kl}^{ij} (respectively $\bar{R}_{kl}^{ij}$), a maximal point $\cap$ labeled by i with the element $-s^{-2i-1}$, and a minimal point $\cup$ labeled by i with the element $-s^{2i+1}$ with $s = \exp\left(\frac{\pi\sqrt{-1}}{N}\right)$ as in Figure 1.




Figure 1

Here R_{kl}^{ij} and $\bar{R}_{kl}^{ij}$ are given by

$$R_{kl}^{ij} = \sum_{n=0}^{\min(N-1-i,j)} \delta_{l,i+n} \delta_{k,j-n} \frac{(i+n)!(N-1+n-j)!}{(i)!(N-1-j)!(n)!} s^{2(i-\frac{N-1}{2})(j-\frac{N-1}{2})-n(i-j)-\frac{n(n+1)}{2}},$$

$$\bar{R}_{kl}^{ij} = \sum_{n=0}^{\min(N-1-j,i)} \delta_{l,i-n} \delta_{k,j+n} \frac{(j+n)!(N-1+n-i)!}{(j)!(N-1-i)!(n)!} (-1)^n$$

$$\times s^{-2(i-\frac{N-1}{2})(j-\frac{N-1}{2})-n(i-j)+\frac{n(n+1)}{2}}$$

with $(n)! = (s - s^{-1})(s^2 - s^{-2}) \cdots (s^n - s^{-n})$.

After multiplying all elements associated to the critical points, we sum up over all labelings. Here we ignore framings of links.

Let us calculate the colored Jones polynomial of the Whitehead link as an example. We can label each edge in the following way, noting Kronecker's deltas in R_{kl}^{ij} and $\bar{R}_{kl}^{ij}$.




Figure 2

We have to rotate a crossing where edges go up. In that case we use $\cup$ and/or $\cap$ to calculate the invariant.

Then we calculate the formula

$$(1) \quad J_N(L) = \sum_{\substack{0 \leq i,j,k \leq N-1 \\ i,j \geq k}} \frac{(q)_i (q)_j \{(q)_{N-1-k}\}^2}{\{(q)_k\}^2 (q)_{N-1-i} (q)_{N-1-j} (q)_{i-k} (q)_{j-k}} q^{-k(i+j+1)},$$

where $q = s^2 = \exp\left(\frac{2\pi\sqrt{-1}}{N}\right)$. Here $(x)_k = (1-x)(1-x^2) \cdots (1-x^k)$.

Next the Chern–Simons invariant of a link is defined as follows. Let $\mathcal{A}$ be the set of all $SO(3)$ -connections of the trivial $SO(3)$ -bundle of a closed three-manifold M and $\text{cs}: \mathcal{A} \rightarrow \mathbb{R}$ the Chern–Simons functional defined by

$$\text{cs}(A) = \frac{1}{8\pi^2} \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right).$$

The Chern–Simons invariant of the connection A is then defined to be the integral

$$\text{cs}_M(A) = \int_{s(M)} \text{cs}(A) \in \mathbb{R}/\mathbb{Z},$$

where the integral is over a section s of the $SO(3)$ -bundle (i.e., an orthonormal frame field on M) [1]. If M is hyperbolic we define $\text{cs}(M)$ to be the Chern–Simons invariant of the connection defined by the hyperbolic metric.

The definition of the Chern–Simons invariant for hyperbolic three-manifolds with cusps is due to R. Meyerhoff [8]. It is defined modulo $1/2$ by using a special singular frame field which is linear near the cusps. See [8] for details. See also [3] how it is computed by SnapPea [12]. Throughout this paper we use another normalization $\text{CS}(M) = -2\pi^2 \text{cs}(M)$ so that $\text{vol}(M) + \sqrt{-1} \text{CS}(M)$ is a natural complexification of the hyperbolic volume $\text{vol}(M)$ (see [10, 15]).

3. KNOT 6_3

Let us calculate the colored Jones polynomial of the knot 6_3 using the labeling as in Figure 3.




Figure 3

Putting $k = n_1 + n_2$ and using the formula in [9]

$$(2) \quad \sum_{i=0}^{N-1} (-1)^i s^{\beta i} \begin{bmatrix} \alpha \\ i \end{bmatrix} = \prod_{j=1}^{\alpha} (1 - s^{\beta + \alpha + 1 - 2j})$$

with $\alpha = k$, $i = n_1$, and $\beta = -k - 1 - 2N$, we calculate

$$J_N(6_3) = \sum_{\substack{0 \leq k, l, m \\ k+l+m \leq N-1}} (-1)^{k+l} s^{\frac{(l+k)(l+k+1)}{2} - \frac{(m+k)(m+k+1)}{2} + \frac{k(k+1)}{2} + 2(m-l)(k+1) + N(m-l+k)} \\ \times \frac{(N-1-l)!(N-1-m)!(l+m+k)!(N-1)!(1-s^{-2N-2}) \cdots (1-s^{-2N-2k})}{(N-1-l-m-k)!(N-1-l-k)!(N-1-m-k)!(l)!(m)!(k)!}.$$

The colored Jones polynomial of the knot 6_3 is given by

$$(3) \quad J_N(6_3) = \sum_{\substack{k, l, m \geq 0 \\ k+l+m \leq N-1}} \left| \frac{(q)_{k+l+m}}{(q)_l (q)_m} \right|^2 (q)_{k+l} (\bar{q})_{m+k} q^{(m-l)(k+1)}.$$

We review of the technique in [6]. For a complex number p and a positive real number γ with $|\operatorname{Re} p| < \pi + \gamma$, we define

$$S_\gamma(p) = \exp \frac{1}{4} \int_{-\infty}^{\infty} \frac{e^{px}}{\sinh(\pi x) \sinh(\gamma x)} \frac{dx}{x}.$$

Here Re denotes the real part. This function has two properties:

- (a) $(1 + \exp(\sqrt{-1}p)) S_\gamma(p + \gamma) = S_\gamma(p - \gamma);$
- (b) $S_\gamma(p) \sim \exp \left(\frac{1}{2\gamma\sqrt{-1}} \operatorname{Li}_2(-\exp(\sqrt{-1}p)) \right) \quad (\gamma \rightarrow 0),$

where

$$\operatorname{Li}_2(z) = - \int_0^z \frac{\log(1-u)}{u} du.$$

We put

$$f_\gamma(p) = \frac{S_\gamma(\gamma - \pi)}{S_\gamma(p)}, \quad \bar{f}_\gamma(p) = \frac{S_\gamma(-p)}{S_\gamma(\pi - \gamma)},$$

so that

$$(q)_k = f_\gamma(-\pi + (2k+1)\gamma), \quad (\bar{q})_k = \bar{f}_\gamma(-\pi + (2k+1)\gamma).$$

Following Kashaev's way, we rewrite the formula (3) as a multiple integral with appropriately chosen contours. (Note that there is considerable doubt as to the contours.) By using the property (b), it can be asymptotically approximated by

$$\iiint \exp \frac{\sqrt{-1}}{2\gamma} V_{6_3}(z, u, v) dz du dv$$

with $\gamma = \pi/N$. Here z , u , and v correspond to q^k , q^m , and q^l respectively, and

$$\begin{aligned} V_{6_3}(z, u, v) = & \operatorname{Li}_2(zuv) - \operatorname{Li}_2\left(\frac{1}{zuv}\right) + \operatorname{Li}_2(zv) - \operatorname{Li}_2\left(\frac{1}{zu}\right) \\ & - \operatorname{Li}_2(u) + \operatorname{Li}_2\left(\frac{1}{u}\right) - \operatorname{Li}_2(v) + \operatorname{Li}_2\left(\frac{1}{v}\right) - \log z \log \frac{u}{v}. \end{aligned}$$

Then there exists a stationary point

$$(z_0, u_0, v_0) = (0.204323 - 0.978904\sqrt{-1}, 1.60838 + 0.558752\sqrt{-1}, 0.554788 + 0.192734\sqrt{-1})$$

of V_{6_3} with

$$\operatorname{Im} V_{6_3}(z_0, u_0, v_0) < 0, \quad \arg z_0 + \arg u_0 + \arg v_0 \leq 2\pi,$$

and we have

$$\begin{aligned} -\operatorname{Im} V_{6_3}(z_0, u_0, v_0) &= 5.693021 \dots, \\ \operatorname{Re} V_{6_3}(z_0, u_0, v_0) &= 0. \end{aligned}$$

From values of $\operatorname{vol}(6_3)$ and $\operatorname{CS}(6_3)$ given by SnapPea, we see that the equation

$$\exp \frac{\sqrt{-1}}{2\gamma} V_{6_3}(z_0, u_0, v_0) = \exp \frac{\operatorname{vol}(6_3) + \sqrt{-1} \operatorname{CS}(6_3)}{2\gamma}$$

holds up to digits shown above.

4. KNOT 8_9

We label the edges of the $(1, 1)$ -tangle presentation of the knot 8_9 as in Figure 4.




Figure 4

We obtain the following formula of the colored Jones polynomial of the knot 8_9 , where we put $l = m_1 + m_2 + k_1 + k_2$ and use the formula (2):

$$J_N(8_9) = \sum_{\substack{0 \leq l, m_1, m_2, n_1, n_2 \leq N-1 \\ m_1 + n_1, m_2 + n_2 \leq l \\ m_1 + m_2 \leq l}} \left| \frac{(q)_{l-m_1} (q)_l (q)_{l-m_2}}{(q)_{m_1} (q)_{m_2} (q)_{n_1} (q)_{n_2}} \right|^2 \frac{(\bar{q})_{l-n_1} (q)_{l-n_2}}{(q)_{l-m_1-n_1} (\bar{q})_{l-m_2-n_2}} \\ \times q^{(m_2-m_1)(l-m_1-m_2) + (n_2-n_1)(l-n_1-n_2) + m_2-m_1+n_2-n_1},$$

which can be asymptotically approximated by

$$\int \cdots \int \exp \frac{\sqrt{-1}}{2\gamma} V_{8_9}(x, y, z, u, v) dx dy dz du dv,$$

where x, y, z, u , and v correspond to $q^{-l}, q^{m_1}, q^{m_2}, q^{n_1}$, and q^{n_2} respectively, and

$$V_{8_9}(x, y, z, u, v) \\ = -\text{Li}_2(xy) + \text{Li}_2\left(\frac{1}{xy}\right) - \text{Li}_2(xz) + \text{Li}_2\left(\frac{1}{xz}\right) - \text{Li}_2(xu) + \text{Li}_2\left(\frac{1}{xv}\right) \\ - \text{Li}_2(x) + \text{Li}_2\left(\frac{1}{x}\right) - \text{Li}_2(y) + \text{Li}_2\left(\frac{1}{y}\right) - \text{Li}_2(z) + \text{Li}_2\left(\frac{1}{z}\right) \\ - \text{Li}_2(u) + \text{Li}_2\left(\frac{1}{u}\right) - \text{Li}_2(v) + \text{Li}_2\left(\frac{1}{v}\right) + \text{Li}_2(xzv) - \text{Li}_2\left(\frac{1}{xyu}\right) \\ - \log \frac{y}{z} \log(xzv) - \log \frac{u}{v} \log(xyu).$$

Consequently we have

$$\begin{aligned} -\operatorname{Im} V_{8_9}(x_0, y_0, z_0, u_0, v_0) &= 7.5881802\dots, \\ \operatorname{Re} V_{8_9}(x_0, y_0, z_0, u_0, v_0) &= 0 \end{aligned}$$

for

$$\begin{aligned} x_0 &= 0.7366011609 - 0.6763273835\sqrt{-1}, \\ y_0 &= 0.4472176075 - 0.1647027124\sqrt{-1}, \\ z_0 &= 1.968989044 - 0.7251455025\sqrt{-1}, \\ u_0 &= 0.3859112582 - 0.0202712198\sqrt{-1}, \\ v_0 &= 2.584139126 - 0.1357401508\sqrt{-1} \end{aligned}$$

satisfying

$$\begin{aligned} \operatorname{Im} V_{8_9}(x_0, y_0, z_0, u_0, v_0) &< 0, \quad \arg x_0 + \arg y_0 + \arg u_0 \leq 2\pi, \\ \arg x_0 + \arg z_0 + \arg v_0 &\leq 2\pi, \quad \arg x_0 + \arg u_0 + \arg v_0 \leq 2\pi. \end{aligned}$$

It follows from the calculation by SnapPea that

$$\exp \frac{\sqrt{-1}}{2\gamma} V_{8_9}(x_0, y_0, z_0, u_0, v_0) = \exp \frac{\operatorname{vol}(8_9) + \sqrt{-1} \operatorname{CS}(8_9)}{2\gamma},$$

up to digits shown above.

5. KNOT 8_{20}

In this section, we discuss a relation between the asymptotic behavior of the colored Jones polynomial and the Chern–Simons invariant for the knot 8_{20} . We label each edge in the diagram of the knot in Figure 5.




Figure 5

The N -colored Jones polynomial of the knot 8_{20} is given by

$$(4) \quad \sum_{\substack{j,l \leq k \leq i+l \leq j+m \\ j \leq i \\ 0 \leq i,j,k,l,m \leq N-1}} \frac{\{(\bar{q})_i(q)_k(\bar{q})_m\}^2}{\{(\bar{q})_j(q)_l\}^2(q)_{k-l}(\bar{q})_{i-k+l}(\bar{q})_{j+m-i-l}(q)_{i-j}(q)_{k-j}} q^{k+m+im+km-il},$$

which can be rewritten in the integral

$$\int \cdots \int \exp \frac{\sqrt{-1}}{2\gamma} V_{8_{20}}(x, y, z, u, v) dx dy dz du dv$$

with

$$\begin{aligned} V_{8_{20}}(x, y, z, u, v) = & -2\text{Li}_2(x) + 2\text{Li}_2\left(\frac{1}{y}\right) + 2\text{Li}_2(z) - 2\text{Li}_2\left(\frac{1}{u}\right) - 2\text{Li}_2\left(\frac{1}{x}\right) \\ & - \text{Li}_2\left(\frac{1}{xy}\right) - \text{Li}_2\left(\frac{z}{y}\right) - \text{Li}_2(zu) + \text{Li}_2(xzu) + \text{Li}_2\left(\frac{1}{xyuv}\right) \\ & + \log x \log u + \log x \log v - \log z \log v + \frac{\pi^2}{2}. \end{aligned}$$

Here x, y, z, u , and v correspond to q^{-i}, q^j, q^k, q^{-l} , and q^m respectively.

Stationary points are solutions to partial differential equations

$$\frac{\partial V_{8_{20}}}{\partial x} = \frac{\partial V_{8_{20}}}{\partial y} = \frac{\partial V_{8_{20}}}{\partial z} = \frac{\partial V_{8_{20}}}{\partial u} = \frac{\partial V_{8_{20}}}{\partial v} = 0.$$

From these equations, we have the following system of algebraic equations

$$\begin{aligned} (1-x)^2 \left(1 - \frac{1}{xyuv}\right) uv &= \left(1 - \frac{1}{xy}\right) (1-xzu), \\ \left(1 - \frac{1}{xy}\right) \left(1 - \frac{z}{y}\right) &= \left(1 - \frac{1}{y}\right)^2 \left(1 - \frac{1}{xyuv}\right), \\ (1-z)^2 (1-xzu) v &= (1-zu) \left(1 - \frac{z}{y}\right), \\ (1-zu) \left(1 - \frac{1}{xyuv}\right) x &= \left(1 - \frac{1}{u}\right)^2 (1-xzu), \\ \left(1 - \frac{1}{v}\right)^2 z &= \left(1 - \frac{1}{xyuv}\right) x. \end{aligned}$$

Using MAPLE V, we get a stationary point $(x_0, y_0, z_0, u_0, v_0)$ which satisfies the conditions

$$\arg \frac{1}{u_0} \leq \arg z_0, \quad \arg z_0 \leq \arg \frac{1}{x_0} + \arg \frac{1}{u_0}$$

from the range in the summation in (4), and

$$\text{Im } V_{8_{20}}(x_0, y_0, z_0, u_0, v_0) < 0,$$

where Im denotes the imaginary part. Note that the range of (4) can be read as

$$\arg \frac{1}{u} \leq \arg z \leq \arg \frac{1}{x} + \arg \frac{1}{u}, \quad \arg \frac{1}{x} + \arg \frac{1}{y} + \arg \frac{1}{u} \leq \arg v,$$

$$0 \leq \arg \frac{1}{x} + \arg \frac{1}{y}, \quad 0 \leq \arg \frac{1}{x}, \arg z, \arg \frac{1}{u}, \arg v \leq 2\pi.$$

To put it concretely,

$$\begin{aligned} x_0 &= 2.878599677 + 2.657408013\sqrt{-1}, \\ y_0 &= \infty, \\ z_0 &= -0.4425377456 - 0.4544788919\sqrt{-1}, \\ u_0 &= 0.3542198353 - 0.02180673815\sqrt{-1}, \\ v_0 &= 0.1458832937 - 0.3399257634\sqrt{-1}. \end{aligned}$$

Then we obtain

$$\begin{aligned} -\operatorname{Im} V_{8_{20}}(x_0, y_0, z_0, u_0, v_0) &= 4.1249032\dots, \\ -\frac{\operatorname{Re} V_{8_{20}}(x_0, y_0, z_0, u_0, v_0) + \pi^2}{2\pi^2} &= 0.1033634\dots \end{aligned}$$

Applying values of $\operatorname{vol}(8_{20})$ and $\operatorname{CS}(8_{20})$ given by SnapPea [12], we see that the following equation holds up to digits shown above.

$$\exp \frac{\sqrt{-1}}{2\gamma} V_{8_{20}}(x_0, y_0, z_0, u_0, v_0) = \exp \frac{\operatorname{vol}(8_{20}) + \sqrt{-1} \operatorname{CS}(8_{20})}{2\gamma}.$$

Note that $\operatorname{CS}(8_{20})$ is defined modulo π^2 .

6. WHITEHEAD LINK

For the final example, we calculate the limit of the colored Jones polynomial of the Whitehead link given by (1), which can be changed to the formula

$$J_N(L) = \sum_{\substack{0 \leq i, j, k \leq N-1 \\ k \leq i, j}} \frac{\{(\bar{q})_i(\bar{q})_j\}^2}{(\bar{q})_k^4(\bar{q})_{i-k}(\bar{q})_{j-k}} q^{-(N-1)N/2}.$$

This can be asymptotically approximated by

$$\iiint \exp \frac{\sqrt{-1}}{2\gamma} V_L(x, y, z) dx dy dz,$$

where

$$V_L(x, y, z) = -2\operatorname{Li}_2\left(\frac{1}{x}\right) - 2\operatorname{Li}_2\left(\frac{1}{y}\right) - 4\operatorname{Li}_2(z) + \operatorname{Li}_2\left(\frac{z}{x}\right) + \operatorname{Li}_2\left(\frac{z}{y}\right) + \pi^2,$$

and x , y , and z correspond to q^i , q^j , and q^k respectively. For a stationary point $(x_0, y_0, z_0) = (\infty, \infty, 1 + \sqrt{-1})$, we obtain

$$\begin{aligned} -\operatorname{Im} V_L(x_0, y_0, z_0) &= 3.663862\dots, \\ -\frac{\operatorname{Re} V_L(x_0, y_0, z_0)}{2\pi^2} &= -0.1250000\dots \end{aligned}$$

Since these values agree with SnapPea, the equation

$$\exp \frac{\sqrt{-1}}{2\gamma} V_L(x_0, y_0, z_0) = \exp \frac{\text{vol}(L) + \sqrt{-1} \text{CS}(L)}{2\gamma}$$

holds up to digits shown above.

7. TOPOLOGICAL CHERN–SIMONS INVARIANT AND SOME EXAMPLES

We propose a topological definition of the Chern–Simons invariant for links.

For a link L , if there exists the limit

$$2\pi \operatorname{Im} \lim_{N \rightarrow \infty} \log \frac{J_{N+1}(L)}{J_N(L)} \pmod{\pi^2},$$

then we denote it by $\text{CS}_{\text{TOP}}(L)$ and call it the *topological Chern–Simons invariant* of L .

Let us give some numerical examples.

For the knot 5_2 , we list some values of $(N, 2\pi \log(J_{N+1}(5_2)/J_N(5_2)))$ by Pari-Gp in the following.

$$\begin{aligned} &(40, 3.058223721261842722613885956 - 3.022924613281720287391974968\sqrt{-1}) \\ &(50, 3.013081508530188353573854822 - 3.023340368517507069134855780\sqrt{-1}) \\ &(60, 2.982744318753580696821772299 - 3.023574042878935429645720640\sqrt{-1}) \\ &(70, 2.960955404961739170749114151 - 3.023717381786374852930574631\sqrt{-1}) \\ &(80, 2.944548269170450112446966301 - 3.023811574968472287718611711\sqrt{-1}) \\ &(100, 2.921483906108228993018469212 - 3.023923719027833555669502480\sqrt{-1}) \\ &(120, 2.906046421388666000282542398 - 3.023985374930307234443986632\sqrt{-1}) \\ &(150, 2.890559881907537128372001511 - 3.024036295143969179028770901\sqrt{-1}) \\ &(200, 2.875024234226941620327156350 - 3.024076266558545340852410631\sqrt{-1}) \\ &(250, 2.865679250969538531562099056 - 3.024094905811349375139149331\sqrt{-1}) \end{aligned}$$

By fitting the above data to quadratic functions on $1/N$, we can obtain the limit value

$$2.82813 - 3.02414\sqrt{-1}$$

of $2\pi \log(J_{N+1}(5_2)/J_N(5_2))$ as $N \rightarrow \infty$ numerically, which agrees with the value

$$2.8281220 - 3.02412837\sqrt{-1}$$

by SnapPea. We display our data graphically in Figure 6 and Figure 7, which help us to see the limit.




Figure 6. Dots indicate $(1/N, 2\pi \operatorname{Re} \log(J_{N+1}(5_2)/J_N(5_2)))$ for $N = 40, 50, 60, 70, 80, 100, 120, 150, 200, 250$. The origin corresponds to $(0, 2.82)$.




Figure 7. Dots indicate $(1/N, 2\pi \operatorname{Im} \log(J_{N+1}(5_2)/J_N(5_2)))$ for $N = 40, 50, 60, 70, 80, 100, 120, 150, 200, 250$. The origin corresponds to $(0, -3.0242)$.

Similarly, for the Whitehead link L , we illustrate our numerical check in Table 1, Figure 8, and Figure 9.

Table 1. $(N, 2\pi \log(J_{N+1}(L)/J_N(L)))$ for the Whitehead link L

(40, 3.892920359101811097809525583 + 2.457483997330866045812504703 $\sqrt{-1}$)
(50, 3.848161466402914225154530180 + 2.461039474018016569869745301 $\sqrt{-1}$)
(60, 3.818029013349499312708236153 + 2.462976748675980254703390855 $\sqrt{-1}$)
(70, 3.796362501209537691078944556 + 2.464147191795881614582476451 $\sqrt{-1}$)
(80, 3.780034327560022195082015385 + 2.464907923404764622274395868 $\sqrt{-1}$)
(100, 3.757062258985477857247991239 + 2.465803785962819679236327339 $\sqrt{-1}$)
(120, 3.741674608179023673159144258 + 2.466291085896660260688606142 $\sqrt{-1}$)
(150, 3.726228649726558590507828429 + 2.466690204011030007962113880 $\sqrt{-1}$)




Figure 8. Dots indicate $(1/N, 2\pi \operatorname{Re} \log(J_{N+1}(L)/J_N(L)))$ for $N = 40, 50, 60, 70, 80, 100, 120, 150$. The origin corresponds to $(0, 3.66)$.




Figure 9. Dots indicate $(1/N, 2\pi \operatorname{Im} \log(J_{N+1}(L)/J_N(L)))$ for $N = 40, 50, 60, 70, 80, 100, 120, 150$. The origin corresponds to $(0, 2.4674)$.

Fitting, we get the numerical limit value $3.66386 + 2.46742\sqrt{-1}$ of $2\pi \log(J_{N+1}(L)/J_N(L))$ as $N \rightarrow \infty$, which agrees with our result in the section 6.

8. CONCLUSION

We have shown the following by concrete calculations.

Observation 8.1. *Let L be one of the hyperbolic knots 6_3 , 8_9 , and 8_{20} , or the Whitehead link. Following Kashaev's way, we approximate the colored Jones polynomial $J_N(L)$ of L asymptotically by*

$$\int \cdots \int \exp \frac{N\sqrt{-1}}{2\pi} V_L(\mathbf{x}) d\mathbf{x}.$$

Then there exists a stationary point $\mathbf{x}_0$ of V_L such that the formula

$$\exp \frac{N\sqrt{-1}}{2\pi} V_L(\mathbf{x}_0) = \exp \frac{N}{2\pi} (\text{vol}(L) + \sqrt{-1} \text{CS}(L))$$

holds up to 6 digits.

Conjecture 8.2 (Complexification of Kashaev's conjecture). *Let L be a hyperbolic link. Then, it holds that*

$$\text{vol}(L) = 2\pi \lim_{N \rightarrow \infty} \frac{\log |\langle L \rangle_N|}{N}$$

with $\text{vol}(L)$ the hyperbolic volume of the complement of L . Moreover, there exists the topological Chern–Simons invariant $\text{CS}_{\text{TOP}}(L)$ of L

$$\text{CS}_{\text{TOP}}(L) = 2\pi \text{Im} \lim_{N \rightarrow \infty} \log \frac{J_{N+1}(L)}{J_N(L)} \mod \pi^2,$$

and $\text{CS}_{\text{TOP}}(L)$ equals to $\text{CS}(L)$ modulo π^2 . Here $\text{CS}(L)$ is the Chern–Simons invariant of L [1, 8]. Note that the complement of L is a hyperbolic manifold with cusps.

We note that Observation 8.1 also holds for the knots 4_1 , 5_2 and 6_1 by calculating Kashaev's examples in [6] using MAPLE V and SnapPea.

Therefore we conclude that the complexified Kashaev's conjecture is true, up to several digits, up to choices of contours when we change summations into integrals, and up to choices of saddle (stationary) points when we approximate integrals by the saddle point method, for the six hyperbolic knots above and for the Whitehead link.

Note that if the complexified Kashaev's conjecture is true then the topological Chern–Simons invariant of a hyperbolic link coincides with its Chern–Simons invariant associated with the hyperbolic metric. Moreover if the volume conjecture is true then the colored Jones polynomial would give both the simplicial volume and the topological Chern–Simons invariant for any knot.

REFERENCES

- [1] S.-S. Chern and J. Simons, *Characteristic forms and geometric invariants*, Ann. of Math. (2) **99** (1974), 48–69. MR 50 #5811
- [2] H. Cohen, *Pari-Gp: a computer program for number theory*, available at <http://www.pari-gp-home.de/>.
- [3] D. Coulson, O.A. Goodman, C.D Hodgson, and W.D. Neumann, *Computing arithmetic invariants of 3-manifolds*, Experiment. Math. **9** (2000), no. 1, 127–152.
- [4] R. M. Kashaev and O. Tirkkonen, *A proof of the volume conjecture on torus knots*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **269** (2000), no. Vopr. Kvant. Teor. Polya i Stat. Fiz. 16, 262–268, 370. MR 1 805 865
- [5] R.M. Kashaev, *A link invariant from quantum dilogarithm*, Modern Phys. Lett. A **10** (1995), no. 19, 1409–1418.
- [6] ———, *The hyperbolic volume of knots from the quantum dilogarithm*, Lett. Math. Phys. **39** (1997), no. 3, 269–275.
- [7] R. Kirby and P. Melvin, *The 3-manifold invariants of Witten and Reshetikhin–Turaev for $sl(2, \mathbb{C})$* , Invent. Math. **105** (1991), 473–545.
- [8] R. Meyerhoff, *Density of the Chern–Simons invariant for hyperbolic 3-manifolds*, Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984), Cambridge Univ. Press, Cambridge, 1986, pp. 217–239. MR **88k**:57033a
- [9] H. Murakami and J. Murakami, *The colored Jones polynomials and the simplicial volume of a knot*, Acta Math. **186** (2001), no. 1, 85–104. MR 1 828 373

- [10] W.D. Neumann and D. Zagier, *Volumes of hyperbolic three-manifolds*, Topology **24** (1985), no. 3, 307–332. MR **87j**:57008
- [11] D. Thurston, *Hyperbolic volume and the Jones polynomial*, Lecture notes, École d'été de Mathématiques 'Invariants de nœuds et de variétés de dimension 3', Institut Fourier - UMR 5582 du CNRS et de l'UJF Grenoble (France) du 21 juin au 9 juillet 1999.
- [12] J. Weeks, *SnapPea: a computer program for creating and studying hyperbolic 3-manifolds*, available at <http://www.northnet.org/weeks/index/SnapPea.html>.
- [13] Y. Yokota, *On the volume conjecture for hyperbolic knots*, preprint, available at http://www.comp.metro-u.ac.jp/~jojo/volume_conjecture.ps.
- [14] ———, *On the volume conjecture of hyperbolic knots*, Knot Theory – dedicated to Professor Kunio Murasugi for his 70th birthday (M. Sakuma, ed.), March 2000, pp. 362–367.
- [15] T. Yoshida, *The η -invariant of hyperbolic 3-manifolds*, Invent. Math. **81** (1985), no. 3, 473–514. MR **87f**:58153

(Hitoshi Murakami) DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, OH-OKAYAMA, MEGURO, TOKYO 152-8551, JAPAN

E-mail address: starshea@tky3.3web.ne.jp

(Jun Murakami) DEPARTMENT OF MATHEMATICAL SCIENCES, SCHOOL OF SCIENCE AND ENGINEERING WASEDA UNIVERSITY 3-4-1, OHKUBO SHINJUKU-KU, TOKYO 169-8555 JAPAN

E-mail address: murakami@mn.waseda.ac.jp

(Miyuki Okamoto) UNIVERSITY OF THE SACRED HEART, HIROO, SHIBUYA, TOKYO 150-8938, JAPAN

E-mail address: miyuki3@hh.iiij4u.or.jp

(Toshie Takata) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, NIIGATA UNIVERSITY, NIIGATA 950-2181, JAPAN

E-mail address: takata@math.sc.niigata-u.ac.jp

(Yoshiyuki Yokota) DEPARTMENT OF MATHEMATICS, TOKYO METROPOLITAN UNIVERSITY, TOKYO 192-0397, JAPAN

E-mail address: jojo@math.metro-u.ac.jp